

## B6.129S1-*Il12a*<sup>tm1Jm</sup>/J

Stock No: 002692

Protocol 29750: Standard PCR Assay - *Il12a*<tm1Jm>

Version 3.2

### Notes

This assay does not work well without the use of a Hotstart Taq polymerase.

The genotyping protocol(s) presented here have been optimized for reagents and conditions used by The Jackson Laboratory (JAX). To genotype animals, JAX recommends researchers validate the assay independently upon receipt of animals into their facility. Reaction cycling temperature and times may require additional optimization based on the specific genotyping reagents used.

### Expected Results

Mutant = 172 bp

Heterozygote = 172 bp and 650 bp

Wild type = 650 bp

Separated by gel electrophoresis on a 1.5% agarose gel.

### JAX Protocol

#### Protocol Primers

PRIMER	5' LABEL	SEQUENCE 5' → 3'	3' LABEL	PRIMER TYPE	REACTION	NOTE
oIMR0158		CTG AAT GAA CTG CAG GAC GA			A	
oIMR0159		ATA CTT TCT CGG CAG GAG CA			A	
oIMR0459		CAG CAT GTG TCA ATC ACG			A	
oIMR0460		TCA CCA TGT CAT CTG TGG			A	

#### Reaction A

COMPONENT	FINAL CONCENTRATION
ddH <sub>2</sub> O	
Kapa 2G HS buffer	1.30 X
MgCl <sub>2</sub>	2.60 mM
dNTP KAPA	0.26 mM
oIMR0158	0.50 uM
oIMR0159	0.50 uM
oIMR0459	0.50 uM
oIMR0460	0.50 uM
Glycerol	6.50 %
Dye	1.00 X
Kapa 2G HS taq polym	0.03 U/ul
DNA	

#### Cycling

STEP	TEMP °C	TIME	NOTE
1	94.0	--	
2	94.0	--	
3	65.0	--	-0.5 C per cycle decrease
4	68.0	--	
5		--	repeat steps 2-4 for 10 cycles (Touchdown)
6	94.0	--	
7	60.0	--	
8	72.0	--	
9		--	repeat steps 6-8 for 28 cycles
10	72.0	--	
11	10.0	--	hold

JAX uses a very high speed Taq (~1000 bp/sec), use cycling times recommended for your reagents.

JAX uses a 'touchdown' cycling protocol and therefore has not calculated the optimal annealing temperature for each set of primers.

