

129-Col4a3^{tm1Dec}/J

Stock No: 002908

Protocol 26326: Separated PCR Assay - Col4a3<tm1Dec>alternate1

Version 1.2

Notes

The genotyping protocol(s) presented here have been optimized for reagents and conditions used by The Jackson Laboratory (JAX). To genotype animals, JAX recommends researchers validate the assay independently upon receipt of animals into their facility. Reaction cycling temperature and times may require additional optimization based on the specific genotyping reagents used.

Expected Results

Mutant = 230 bp

Heterozygote = 230 bp and 309 bp

Wild type = 309 bp

JAX Protocol

Protocol Primers

PRIMER	5' LABEL	SEQUENCE 5' → 3'	3' LABEL	PRIMER TYPE	REACTION	NOTE
14885		TGC TCT CTC AAA TGC ACC AG		Wild type Reverse	B	
oIMR0643		CCA GGC TTA AAG GGA AAT CC		Common	A, B	
oIMR1100		GCT ATC AGG ACA TAG CGT TGG		Mutant Reverse	A	

Reaction A

COMPONENT	FINAL CONCENTRATION
ddH ₂ O	
Kapa 2G HS buffer	1.30 X
MgCl ₂	2.60 mM
dNTPS-kapa	0.26 mM
oIMR0643	0.50 uM
oIMR1100	0.50 uM
Glycerol	6.50 %
Dye	1.00 X
Kapa 2G HS taq polym	0.03 U/ul
DNA	

Cycling

STEP	TEMP °C	TIME	NOTE
1	94.0	--	
2	94.0	--	
3	65.0	--	-0.5 C per cycle decrease
4	68.0	--	
5		--	repeat steps 2-4 for 10 cycles (Touchdown)
6	94.0	--	
7	60.0	--	
8	72.0	--	
9		--	repeat steps 6-8 for 28 cycles
10	72.0	--	
11	10.0	--	hold

JAX uses a very high speed Taq (~1000 bp/sec), use cycling times recommended for your reagents.

JAX uses a 'touchdown' cycling protocol and therefore has not calculated the optimal annealing temperature for each set of primers.

Reaction B

COMPONENT	FINAL CONCENTRATION
ddH2O	
Kapa 2G HS buffer	1.30 X
MgCl2	2.60 mM
dNTPS-kapa	0.26 mM
14885	0.50 uM
oIMR0643	0.50 uM
Glycerol	6.50 %
Dye	1.00 X
Kapa 2G HS taq polym	0.03 U/ul
DNA	

Cycling

STEP	TEMP °C	TIME	NOTE
1	94.0	--	
2	94.0	--	
3	65.0	--	-0.5 C per cycle decrease
4	68.0	--	
5		--	repeat steps 2-4 for 10 cycles (Touchdown)
6	94.0	--	
7	60.0	--	
8	72.0	--	
9		--	repeat steps 6-8 for 28 cycles
10	72.0	--	
11	10.0	--	hold

JAX uses a very high speed Taq (~1000 bp/sec), use cycling times recommended for your reagents.

JAX uses a 'touchdown' cycling protocol and therefore has not calculated the optimal annealing temperature for each set of primers.

