

STOCK Tg(CAG-Bgeo/ALPP)1Lbe/J

Stock No: 003919

Protocol 22297: Standard PCR Assay - Tg(ALPP)

Version 1.3

Notes

The genotyping protocol(s) presented here have been optimized for reagents and conditions used by The Jackson Laboratory (JAX). To genotype animals, JAX recommends researchers validate the assay independently upon receipt of animals into their facility. Reaction cycling temperature and times may require additional optimization based on the specific genotyping reagents used.

Expected Results

TG = 548 bp

internal control = 200 bp

Separated by gel electrophoresis on a 1.5% agarose gel.

JAX Protocol

Protocol Primers

PRIMER	5' LABEL	SEQUENCE 5' → 3'	3' LABEL	PRIMER TYPE	REACTION	NOTE
oIMR1382		CCG CTT CCC ATA TGT GGC TCT GTC C		Transgene	A	
oIMR1383		GCA TGA GCT CAG TGC GGT TCC ACA C		Transgene	A	
oIMR8744		CAA ATG TTG CTT GTC TGG TG		Internal Positive Control Forward	A	
oIMR8745		GTC AGT CGA GTG CAC AGT TT		Internal Positive Control Reverse	A	

Reaction A

COMPONENT	FINAL CONCENTRATION
ddH ₂ O	
Kapa 2G HS buffer	1.30 X
MgCl ₂	2.60 mM
dNTP KAPA	0.26 mM
oIMR1382	0.50 uM
oIMR1383	0.50 uM
oIMR8744	0.50 uM
oIMR8745	0.50 uM
Glycerol	6.50 %
Dye	1.00 X
Kapa 2G HS taq polym	0.03 U/ul
DNA	

Cycling

STEP	TEMP °C	TIME	NOTE
1	94.0	--	
2	94.0	--	
3	65.0	--	-0.5 C per cycle decrease
4	68.0	--	
5		--	repeat steps 2-4 for 10 cycles (Touchdown)
6	94.0	--	
7	60.0	--	
8	72.0	--	
9		--	repeat steps 6-8 for 28 cycles
10	72.0	--	
11	10.0	--	hold

JAX uses a very high speed Taq (~1000 bp/sec), use cycling times recommended for your reagents.

JAX uses a 'touchdown' cycling protocol and therefore has not calculated the optimal annealing temperature for each set of primers.

