

NOD.129S2(B6)-*Casp1*^{tm1Sesh} *Casp4*^{del}/LtJ

Stock No: 004947

Protocol 27941: Separated PCR Assay - Casp1<tm1Sesh>

Version 5.2

Notes

The genotyping protocol(s) presented here have been optimized for reagents and conditions used by The Jackson Laboratory (JAX). To genotype animals, JAX recommends researchers validate the assay independently upon receipt of animals into their facility. Reaction cycling temperature and times may require additional optimization based on the specific genotyping reagents used.

Expected Results

Mutant = 600bp

Wild type = 600bp

Separated by gel electrophoresis on a 1.5% agarose gel. Do not pool as both bands are approximately

Sequence

Run Wt with LR and no dye

JAX Protocol

Protocol Primers

| PRIMER | 5' LABEL | SEQUENCE 5' → 3' | 3' LABEL | PRIMER TYPE | REACTION | NOTE |
|----------|----------|-----------------------------|----------|-------------|----------|------|
| oIMR3769 | | GAA GAG ATG TTA CAG AAG CC | | | A, B | |
| oIMR3770 | | CAT GCC TGA ATA ATG ATC ACC | | | A | |
| oIMR3771 | | GCG CCT CCC CTA CCC GG | | | B | |

Reaction A

| COMPONENT | FINAL CONCENTRATION |
|----------------------|---------------------|
| ddH2O | |
| Kapa 2G HS buffer | 1.30 X |
| MgCl ₂ | 2.60 mM |
| dNTPS-kapa | 0.26 mM |
| oIMR3769 | 0.50 uM |
| oIMR3770 | 0.50 uM |
| Glycerol | 6.50 % |
| Dye | 1.00 X |
| Kapa 2G HS taq polym | 0.03 U/ul |
| DNA | |

Cycling

| STEP | TEMP °C | TIME | NOTE |
|------|---------|------|--|
| 1 | 94.0 | -- | |
| 2 | 94.0 | -- | |
| 3 | 65.0 | -- | -0.5 C per cycle decrease |
| 4 | 68.0 | -- | |
| 5 | | -- | repeat steps 2-4 for 10 cycles (Touchdown) |
| 6 | 94.0 | -- | |
| 7 | 60.0 | -- | |
| 8 | 72.0 | -- | |
| 9 | | -- | repeat steps 6-8 for 28 cycles |
| 10 | 72.0 | -- | |
| 11 | 10.0 | -- | hold |

JAX uses a very high speed Taq (~1000 bp/sec), use cycling times recommended for your reagents.

JAX uses a 'touchdown' cycling protocol and therefore has not calculated the optimal annealing temperature for each set of primers.

Reaction B

| COMPONENT | FINAL CONCENTRATION |
|----------------------|---------------------|
| ddH2O | |
| Kapa 2G HS buffer | 1.30 X |
| MgCl2 | 2.60 mM |
| dNTPS-kapa | 0.26 mM |
| oIMR3769 | 0.50 uM |
| oIMR3771 | 0.50 uM |
| Glycerol | 6.50 % |
| Dye | 1.00 X |
| Kapa 2G HS taq polym | 0.03 U/ul |
| DNA | |

Cycling

| STEP | TEMP °C | TIME | NOTE |
|------|---------|------|--|
| 1 | 94.0 | -- | |
| 2 | 94.0 | -- | |
| 3 | 65.0 | -- | -0.5 C per cycle decrease |
| 4 | 68.0 | -- | |
| 5 | | -- | repeat steps 2-4 for 10 cycles (Touchdown) |
| 6 | 94.0 | -- | |
| 7 | 60.0 | -- | |
| 8 | 72.0 | -- | |
| 9 | | -- | repeat steps 6-8 for 28 cycles |
| 10 | 72.0 | -- | |
| 11 | 10.0 | -- | hold |

JAX uses a very high speed Taq (~1000 bp/sec), use cycling times recommended for your reagents.

JAX uses a 'touchdown' cycling protocol and therefore has not calculated the optimal annealing temperature for each set of primers.

