

## B6;129-Slc30a3<sup>tm1Rpa</sup>/J

Stock No: 005064

Protocol 33859: Separated PCR Assay - Slc30a3<tm1Rpa>

Version 2.0

### Notes

The genotyping protocol(s) presented here have been optimized for reagents and conditions used by The Jackson Laboratory (JAX). To genotype animals, JAX recommends researchers validate the assay independently upon receipt of animals into their facility. Reaction cycling temperature and times may require additional optimization based on the specific genotyping reagents used.

### Expected Results

Mutant = 400 bp

Heterozygote = 650 bp and 400 bp

Wild type = 650 bp

Separated by gel electrophoresis on a 1.5% agarose gel.

### JAX Protocol

#### Protocol Primers

| PRIMER   | 5' LABEL | SEQUENCE 5' → 3'               | 3' LABEL | PRIMER TYPE | REACTION | NOTE |
|----------|----------|--------------------------------|----------|-------------|----------|------|
| oIMR3663 |          | CCT GTG CTC TAG TAG CTT TAC GG |          | Mutant      | B        |      |
| oIMR3693 |          | AGT CAC TGG CAT CCT CCT GT     |          | Wild type   |          |      |
| oIMR3694 |          | GGT ATC CAT GCC CTT CCT CT     |          | Common      |          |      |

#### Reaction A

| COMPONENT            | FINAL CONCENTRATION |
|----------------------|---------------------|
| ddH <sub>2</sub> O   |                     |
| Kapa 2G HS buffer    | 1.30 X              |
| MgCl <sub>2</sub>    | 2.60 mM             |
| dNTPS-kapa           | 0.26 mM             |
| Primer 1             | 0.50 uM             |
| Primer 2             | 0.50 uM             |
| Glycerol             | 6.50 %              |
| Dye                  | 1.00 X              |
| Kapa 2G HS taq polym | 0.03 U/ul           |
| DNA                  |                     |

#### Cycling

| STEP | TEMP °C | TIME | NOTE                                       |
|------|---------|------|--|
| 1    | 94.0    | --   |  |
| 2    | 94.0    | --   |  |
| 3    | 65.0    | --   | -0.5 C per cycle decrease                  |
| 4    | 68.0    | --   |  |
| 5    |         | --   | repeat steps 2-4 for 10 cycles (Touchdown) |
| 6    | 94.0    | --   |  |
| 7    | 60.0    | --   |  |
| 8    | 72.0    | --   |  |
| 9    |         | --   | repeat steps 6-8 for 28 cycles             |
| 10   | 72.0    | --   |  |
| 11   | 10.0    | --   | hold                                       |

JAX uses a very high speed Taq (~1000 bp/sec), use cycling times recommended for your reagents.

JAX uses a 'touchdown' cycling protocol and therefore has not calculated the optimal annealing temperature for each set of primers.

## Reaction B

| COMPONENT            | FINAL CONCENTRATION |
|----------------------|---------------------|
| ddH2O                |                     |
| Kapa 2G HS buffer    | 1.30 X              |
| MgCl2                | 2.60 mM             |
| dNTPS-kapa           | 0.26 mM             |
| oIMR3663             | 0.50 uM             |
| Primer 2             | 0.50 uM             |
| Glycerol             | 6.50 %              |
| Dye                  | 1.00 X              |
| Kapa 2G HS taq polym | 0.03 U/ul           |
| DNA                  |                     |

## Cycling

| STEP | TEMP °C | TIME | NOTE                                       |
|------|---------|------|--|
| 1    | 94.0    | --   |  |
| 2    | 94.0    | --   |  |
| 3    | 65.0    | --   | -0.5 C per cycle decrease                  |
| 4    | 68.0    | --   |  |
| 5    |         | --   | repeat steps 2-4 for 10 cycles (Touchdown) |
| 6    | 94.0    | --   |  |
| 7    | 60.0    | --   |  |
| 8    | 72.0    | --   |  |
| 9    |         | --   | repeat steps 6-8 for 28 cycles             |
| 10   | 72.0    | --   |  |
| 11   | 10.0    | --   | hold                                       |

JAX uses a very high speed Taq (~1000 bp/sec), use cycling times recommended for your reagents.

JAX uses a 'touchdown' cycling protocol and therefore has not calculated the optimal annealing temperature for each set of primers.

