

## STOCK *Hhat*<sup>Tg(TFAP2A-cre)</sup>1Will/J

Stock No: 023407

Protocol 27144: Standard PCR Assay - Hhat<Tg(TFAP2A-cre)1Will>

Version 1.2

### Notes

The genotyping protocol(s) presented here have been optimized for reagents and conditions used by The Jackson Laboratory (JAX). To genotype animals, JAX recommends researchers validate the assay independently upon receipt of animals into their facility. Reaction cycling temperature and times may require additional optimization based on the specific genotyping reagents used.

### Expected Results

Mutant = ~275 bp

Heterozygote = ~275 bp and 699 bp

Wild type = 699 bp

### JAX Protocol

#### Protocol Primers

PRIMER	5' LABEL	SEQUENCE 5' → 3'	3' LABEL	PRIMER TYPE	REACTION	NOTE
20277		GCC TGT AAA GTC ACA TGC TCT G		Common	A	
20278		TCT CAT CTG GCC TGT CCT CT		Wild type Reverse	A	
20279		CAC TTG CTA ACT AGA AGG AAC TTC C		Transgene Reverse	A	

#### Reaction A

COMPONENT	FINAL CONCENTRATION
ddH <sub>2</sub> O	
Kapa 2G HS buffer	1.30 X
MgCl <sub>2</sub>	2.60 mM
dNTP KAPA	0.26 mM
20277	0.50 uM
20278	0.50 uM
20279	0.50 uM
Glycerol	6.50 %
Dye	1.00 X
Kapa 2G HS taq polym	0.03 U/ul
DNA	

#### Cycling

STEP	TEMP °C	TIME	NOTE
1	94.0	--	
2	94.0	--	
3	65.0	--	-0.5 C per cycle decrease
4	68.0	--	
5		--	repeat steps 2-4 for 10 cycles (Touchdown)
6	94.0	--	
7	60.0	--	
8	72.0	--	
9		--	repeat steps 6-8 for 28 cycles
10	72.0	--	
11	10.0	--	hold

JAX uses a very high speed Taq (~1000 bp/sec), use cycling times recommended for your reagents.

JAX uses a 'touchdown' cycling protocol and therefore has not calculated the optimal annealing temperature for each set of primers.

